arXiv:1111.1385v3 [math.GR] 22 Aug 2012

Two approaches towards property (T) for groups acting on simplicial complexes

Izhar Oppenheim

Department of Mathematics, The Technion-Israel Institute of Technology, 32000 Haifa, Israel
E-mail: izharo@gmail.com

August 24, 2012

Abstract. This article generalizes two approaches for property (T) - the first is a generalization of Żuk's criterion for property (T) and the second is a generalization of the work of Kassabov regarding property (T) and subspace arrangements. In both cases we obtain new criteria for property (T) (and for vanishing of higher L^2-cohomologies).

Mathematics Subject Classification (2010). Primary 20F65; Secondary 05C25

Keywords. Property (T), cohomology, simplicial complex, Laplacian

1 Introduction

We'll start by briefly discussing two approaches toward proving property (T).

1.1 Criteria for property (T) regarding Laplacian eigenvalues

For a finite graph L with a set of vertices V_L, the Laplacian of the graph Δ^+ is an operator on the space of real valued functions on V_L which is defined as

$$\Delta^+ f(v) = f(v) - \frac{1}{m(v)} \sum_{u \sim v} f(u)$$

where $m(v)$ is the valance of v and $u \sim v$ means that there is an edge connecting u and v. The Laplacian is a positive operator and we denote by $\lambda(L)$ its smallest positive eigenvalue. One can generalize the definition of the Laplacian so it will be defined for a simplicial complex X of any dimension. For such a complex the

Laplacian is again a positive operator and we denote by $\lambda(X)$ its smallest positive eigenvalue.

Ballmann and Światkowski in [BŚ97] and Żuk in [Żuk96] gave criteria for the vanishing of the L^2-cohomology of a group Γ acting on a simplicial complex X, by considering the values of λ for the links of X. More in specifically [BŚ97] and [Żuk96] proved the following theorems:

[BŚ97, Theorem 2.5]. *Let X be a locally finite simplicial complex of dimension n and let Γ be a group acting properly discontinuously and by automorphisms on X. Assume that for every simplex η of dimension $k-1$, the link of η denoted by X_τ is a connected simplicial complex and that there is $\varepsilon > 0$ such that $\lambda(X_\eta) \geq \frac{k(n-k)}{k+1} + \varepsilon$, then $L^2H^k(X,\rho) = 0$ for any unitary representation ρ of Γ.*

[Żuk96, Theorem 1]. *Let X be a locally finite simplicial complex of dimension 2 and let Γ be a group acting properly discontinuously and freely by automorphisms on X. Assume that for every vertex v, the link of v denoted by X_v is a connected graph and that for every two connected vertices u, v in X we have the following inequality $\frac{\lambda(X_u)+\lambda(X_v)}{2} > \frac{1}{2}$, then $L^2H^1(X,\rho) = 0$ for any unitary representation ρ of Γ.*

In [DJ00] the above theorems were generalized by Dymara and Januszkiewicz to a more general setting in which Γ isn't necessarily discrete but just locally compact and unimodular.

If Γ acts cocompactly and properly discontinuously by automorphisms on a contractible simplicial complex X, then the following are equivalent:

1. Γ has property (T).
2. $L^2H^1(X,\rho) = 0$ for any unitary representation ρ of Γ (see [BdlHV08]).
3. $L^2\mathcal{H}^1(X,\rho) = 0$ for any unitary representation ρ of Γ, where $L^2\mathcal{H}^1(X,\rho)$ is the first reduced L^2-cohomology (see [BdlHV08], [Sha00]).

Therefore the above theorems give criteria for property (T) when Γ acts cocompactly and properly on a contractible locally finite simplicial complex of dimension 2.

1.2 Criterion for property (T) regarding angles

A different approach towards property (T) was taken in by Kassabov in [Kas11]. In [Kas11] a concept of a cosine, denote as $cos(V,U)$, between two Hilbert subspaces V, U is defined. This concept is used to in the following theorem:

[Kas11, Theorem 5.1]. *Let $V_1, ..., V_n$ be closed subspaces of a Hilbert space H. Suppose that the $n \times n$ symmetric matrix*

$$A = \begin{pmatrix} 1 & -\varepsilon_{12} & -\varepsilon_{13} & \cdots & -\varepsilon_{1n} \\ -\varepsilon_{21} & 1 & -\varepsilon_{23} & \cdots & -\varepsilon_{2n} \\ -\varepsilon_{31} & -\varepsilon_{32} & 1 & \cdots & -\varepsilon_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\varepsilon_{n1} & -\varepsilon_{n2} & -\varepsilon_{n3} & \cdots & 1 \end{pmatrix}$$

where $\varepsilon_{ij} = cos(V_i, V_j)$ *is positive definite. Then for every* $v \in H$ *we have*

$$d(v, \bigcap V_i) \leq d_v^t A^{-1} d_v$$

where d_v *is the column vector with entries* $d(v, V_i)$*.*

This theorem can be used to give a criterion for property (T) in the case that $\Gamma = \langle G_1, ..., G_n \rangle$, where $G_1, ..., G_n$ are compact groups. The criterion is as follows: for a unitary representation $\rho : \Gamma \to \mathcal{U}(H_\rho)$ denote by $H_\rho^{G_i}$ the closed subspace of H_ρ fixed by G_i. Define

$$\varepsilon_{ij} = inf_\rho \{cos(H_\rho^{G_i}, H_\rho^{G_j})\}$$

if

$$A = \begin{pmatrix} 1 & -\varepsilon_{12} & -\varepsilon_{13} & \cdots & -\varepsilon_{1n} \\ -\varepsilon_{21} & 1 & -\varepsilon_{23} & \cdots & -\varepsilon_{2n} \\ -\varepsilon_{31} & -\varepsilon_{32} & 1 & \cdots & -\varepsilon_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\varepsilon_{n1} & -\varepsilon_{n2} & -\varepsilon_{n3} & \cdots & 1 \end{pmatrix}$$

is a positive matrix, then Γ has property (T) (we refer the reader to [Kas11, Observation 2.1] for the explanation of the connection between the theorem and the criterion).

Observe that this theorem can be applied in the case that Γ is a group acting properly on an n-dimensional connected simplicial complex and the fundamental domain is a single (n-dimensional) simplex denoted by σ. In this case, Γ is generated by the stabilizers of the $(n-1)$-dimensional faces of σ denoted by $G_1, ..., G_{n+1}$. Note that since the action is proper, we get that G_i is compact for every i.

1.3 New criteria for property (T)

In this article we shall generalize the above approaches to get new criteria for vanishing of L^2-cohomologies for groups acting on simplicial complexes. The two generalizations are of different nature: the generalization of the [BŚ97, Theorem 2.5] and [Żuk96, Theorem 1] are basically some sort of averaging along a simplex of the first positive eigenvalues of the Laplacian at the links. For the 2-dimensional case we get the following theorem:

Theorem 1. *Let* Γ *be a locally compact, properly discontinuous, unimodular group of automorphisms of* X *acting cocompactly on* X *such that* X *is a locally finite contractible* 2*-dimensional simplicial complex. For a* 1*-simplex* (u, v) *of* X *denote*

$$S_{(u,v)} = \lambda(X_u) + \lambda(X_v) - 1$$

If for every 2*-simplex* (u, v, w) *of* X *the following holds:*

- $\lambda(X_u) + \lambda(X_v) + \lambda(X_w) > \frac{3}{2}$
- $S_{(u,v)} S_{(u,w)} + S_{(u,v)} S_{(v,w)} + S_{(u,w)} S_{(v,w)} > 0$

then Γ *has property (T).*

There are some interesting examples for the above theorem coming from the sporadic finite simple groups. Here is one such example: in [Kan81] it is shown that the Lyons group (which is a sporadic finite simple group) acts on a finite simplicial complex Y and that the universal cover of Y is an exotic affine building. This yields an extension of the Lyons group by $\pi_1(Y)$:

$$0 \longrightarrow \pi_1(Y) \longrightarrow \widetilde{Lyons} \longrightarrow Lyons \longrightarrow 0$$

As it turns out, Żuk's criterion doesn't show that $\pi_1(Y)$ has property (T) but our criterion holds and we are able to prove property (T) for $\pi_1(Y)$ and $\widetilde{Lyons}$. There are 2 other examples of this nature in the last section of this article. In addition, there are some hyperbolic Kac-Moody groups where our criterion holds and Żuk's criterion fails.

The generalization of [Kas11, Theorem 5.1] deals with the case that the fundamental domain of the group action is not a single simplex (as in [Kas11]) but a finite simplicial complex. In order to state our theorem we introduce the following terminology: let $Y = (V, E)$ be a connected finite graph (with no loops or multiple edges), Λ be a subgroup of $Aut(Y)$ and $\rho : \Lambda \to \mathcal{U}(H)$ be a unitary representation (H is a Hilbert space). Denote

$$C^0(Y, \rho) = \{\phi : V \to H : \phi \text{ is equivariant}\}$$

Define the reduced cosine of Y with respect to Γ and ρ to be

$$cos_r(Y, \Lambda, \rho) = sup\{\frac{\sum_{(u,v)\in E}\langle \phi(u), \phi(v)\rangle}{\sum_{(u,v)\in E} |\phi(u)||\phi(v)|} : 0 \neq \phi \in C^0(Y, \rho), \sum_{(u,v)\in E} \phi(u)+\phi(v) = 0\}$$

and define

$$cos_r(Y, \Lambda) = sup\{cos_r(Y, \Gamma, \rho) : \rho \text{ is a unitary representation of } \Gamma\}$$

Now we can state the theorem for in the 2-dimensional case:

Theorem 2. *Let* Γ *be a locally compact, properly discontinuous, unimodular group of automorphisms of* X *acting cocompactly on* X *such that* X *is a locally finite contractible* 2*-dimensional simplicial complex. For a vertex* u *denote by* Γ_u *the stabilizer of* u *in* Γ *acting on* X_u*. If for every* 2*-simplex* (u, v, w) *of* X *the matrix:*

$$A(u, v, w) = \begin{pmatrix} 1 & -cos_r(X_u, \Gamma_u) & -cos_r(X_v, \Gamma_v) \\ -cos_r(X_u, \Gamma_u) & 1 & -cos_r(X_w, \Gamma_w) \\ -cos_r(X_v, \Gamma_v) & -cos_r(X_w, \Gamma_w) & 1 \end{pmatrix}$$

is positive definite then Γ *has property (T).*

Structure of the paper. Section 2 is devoted to introducing the framework developed in [BŚ97] and [DJ00], section 3 contains the vanishing results for L^2-cohomologies relaying on Laplacian eigenvalue, section 4 contains vanishing results for L^2-cohomologies relaying on cosines of links and section 5 contains examples of groups in which Theorem 1 proves property (T) (sadly, at the this time we are not able to produce examples for Theorem 2 that differ from the ones given in [Kas11]) .

Acknowledgments. The author wishes to thank his advisor Uri Bader, for many useful conversations and much encouragement along the way.

2 Framework

Here we introduce the framework constructed in [BŚ97] and [DJ00]. Throughout this section X is a locally finite simplicial complex of dimension n such that all the links of X are connected, Γ is a locally compact, properly discontinuous, unimodular group of automorphisms of X and ρ is a unitary representation of Γ on a complex Hilbert space H.

2.1 general settings

Following [BŚ97] we introduce the following notations:

1. For $0 \leq k \leq n$, denote by $\Sigma(k)$ the set of ordered k-simplices (i.e. $\sigma \in \Sigma(k)$ is and ordered $k+1$-tuple of vertices) and choose a set $\Sigma(k, \Gamma) \subseteq \Sigma(k)$ of representatives of Γ-orbits.
2. For a simplex $\sigma \in \Sigma(k)$ denote by $m(\sigma)$ the number of n-simplices containing σ without ordering, i.e. if $\sigma = (v_0, ..., v_k)$ then $m(\sigma)$ is the number of different unordered n-simplices which contain $\{v_0, ..., v_k\}$. We shall assume that $m(\sigma) \geq 1$ for every σ.
3. For $\tau = (w_0, ..., w_k), \sigma = (v_0, ..., v_l)$ s.t. $k \leq l$, we denote $\tau \subset \sigma$ if $\{w_0, ..., w_k\} \subset \{v_0, ..., v_l\}$ and $\tau \sqsubset \sigma$ if there is a monotone function $f : \{0, ..., k\} \rightarrow \{0, ..., l\}$ such that $w_i = v_{f(i)}$.
4. For a simplex $\sigma \in \Sigma(k)$, denote by Γ_σ the stabilizer of σ and by $|\Gamma_\sigma|$ the measure of Γ_σ with respect to the Haar measure.
5. For $0 \leq k \leq n$, denote by $C^k(X, \rho)$ the space of simplicial k-cochains of X which are twisted by ρ, that is, $\phi \in C^k(X, \rho)$ is an alternating map on ordered k-simplices of X with values in H such that

$$\forall \gamma \in \Gamma, \forall x \in \Sigma(k), \rho(\gamma)\phi(x) = \phi(\gamma x)$$

For $\phi \in C^k(X, \rho)$ define ϕ to be *square integrable mod* Γ if

$$\|\phi\|^2 := \sum_{\sigma \in \Sigma(k,\Gamma)} \frac{m(\sigma)}{(k+1)! \, |\Gamma_\sigma|} \langle \phi(\sigma), \phi(\sigma) \rangle < \infty$$

Denote by $L^k(X,\rho)$ the space of all square integrable cochains in $C^k(X,\rho)$. On $L^k(X,\rho)$ there is an Hermitian form given by

$$\langle \phi, \psi \rangle := \sum_{\sigma \in \Sigma(k,\Gamma)} \frac{m(\sigma)}{(k+1)!\,|\Gamma_\sigma|} \langle \phi(\sigma), \psi(\sigma) \rangle \,,\ \phi, \psi \in L^k(X,\rho)$$

Note that if Γ acts cocompactly on X then $L^k(X,\rho) = C^k(X,\rho)$. To distinguish the norm of $L^k(X,\rho)$ from the norm of H we will use $|.|$ for the norm of H (i.e. $\langle \phi(\sigma), \phi(\sigma) \rangle = |\phi(\sigma)|^2$).

For $0 \leq k < n$, the *differential* $d : C^k(X,\rho) \to C^{k+1}(X,\rho)$ is given by

$$d\phi(\sigma) := \sum_{i=0}^{k+1} (-1)^i \phi(\sigma_i),\ \sigma \in \Sigma(k+1)$$

Where $\sigma_i = (v_0, ..., \hat{v_i}, ..., v_{k+1})$ for $(v_0, ..., v_{k+1}) = \sigma \in \Sigma(k+1)$.

Proposition 2.1. *In [BŚ97] the following facts are proved:*

1. *[BŚ97, Proposition 1.5] When the differential is restricted to* $d : L^k(X,\rho) \to L^{k+1}(X,\rho)$ *it is a bounded operator.*
2. *[BŚ97, Proposition 1.6] The adjoint operator of* d*, denoted by* $\delta : L^{k+1}(X,\rho) \to L^k(X,\rho)$ *is*

$$\delta\phi(\tau) = \sum_{v \in \Sigma(0), v\tau \in \Sigma(k+1)} \frac{m(v\tau)}{m(\tau)} \phi(v\tau),\ \tau \in \Sigma(k)$$

   where $v\tau = (v, v_0, ..., v_k)$ *for* $\tau = (v_0, ..., v_k)$

3. *[BŚ97, Corollary 1.7] For* $\phi \in L^k(X,\rho)$ *and* $\sigma \in \Sigma(k)$*,*

$$\delta d\phi(\sigma) = (n-k)\phi(\sigma) - \sum_{v \in \Sigma(0), v\tau \in \Sigma(k+1)} \sum_{0 \leq i \leq k} (-1)^i \frac{m(v\sigma)}{m(\sigma)} \phi(v\sigma_i)$$

Remark 2.2. *Note that if* X *is a 1-dimentional connected graph,* $k = 0$ *and* $H = \mathbb{R}$*, then* δd *is the (classical) graph Laplacian.*

2.2 L^2-cohomology

The L^2-cohomology of X with respect to ρ is defined as

$$L^2H^k(X,\rho) = ker(d \mid L^k(X,\rho)) / im(d \mid L^{k-1}(X,\rho))$$

And the reduced L^2-cohomology of X with respect to ρ is defined as

$$ker(d \mid L^k(X,\rho)) / \overline{im(d \mid L^{k-1}(X,\rho))}$$

Note that if Γ acts cocompactly on a contractible X then $L^2H^k(X,\rho) = L^2H^k(\Gamma,\rho)$.

Let

$$\Delta^+ = \delta d, \Delta^- = d\delta, \Delta = \Delta^+ + \Delta^-$$

Define the space of $L^2\mathcal{H}^k(X,\rho)$ of square integrable mod Γ *harmonic k-forms* twisted by ρ to be

$$L^2\mathcal{H}^k(X,\rho) = ker(\Delta \mid L^k(X,\rho)) = ker(\Delta^+ \mid L^k(X,\rho)) \cap ker(\Delta^- \mid L^k(X,\rho))$$

We have

$$(ker(\Delta^+ \mid L^k(X,\rho)))^\perp = \overline{im(\Delta^+ \mid L^k(X,\rho))} = \overline{im(\delta \mid L^{k+1}(X,\rho))}$$

$$(ker(\Delta^- \mid L^k(X,\rho)))^\perp = \overline{im(\Delta^- \mid L^k(X,\rho))} = \overline{im(d \mid L^{k-1}(X,\rho))}$$

and the orthogonal decompositions

$$ker(\Delta^+ \mid L^k(X,\rho)) = L^2\mathcal{H}^k(X,\rho) \oplus \overline{im(\Delta^- \mid L^k(X,\rho))}$$

$$ker(\Delta^- \mid L^k(X,\rho)) = L^2\mathcal{H}^k(X,\rho) \oplus \overline{im(\Delta^+ \mid L^k(X,\rho))}$$

In particular,

$$L^2\mathcal{H}^k(X,\rho) = ker(d \mid L^k(X,\rho))/\overline{im(d \mid L^{k-1}(X,\rho))}$$

So $L^2\mathcal{H}^k(X,\rho)$ equals the reduced cohomology.

2.3 Localization

Let $(v_0,...,v_j) = \tau \in \Sigma(j)$, denote by X_τ the *link* of τ in X, that is, the subcomplex of dimension $n-j-1$ consisting on simplices $\sigma = (w_0,...,w_k)$ such that $\{v_0,...,v_j\}, \{w_0,...,w_k\}$ are disjoint as sets and $(v_0,...,v_j,w_0,...,w_k) = \tau\sigma \in \Sigma(j+k+1)$. The isotropy group Γ_τ acts by automorphisms on X_τ and if we denote by ρ_τ the restriction of ρ to Γ_τ, we get that ρ_τ is a unitary representation of Γ_τ. Note that since X is locally finite and Γ acts properly discontinuously, we get that X_τ is finite and Γ_τ is compact. Denote as in the general settings:

1. For $0 \leq k \leq n-j-1$, denote by $\Sigma_\tau(k)$ the set of ordered k-simplices and choose a set $\Sigma_\tau(k,\Gamma_\tau) \subseteq \Sigma_\tau(k)$ of representatives of Γ_τ-orbits.
2. For a simplex $\sigma \in \Sigma_\tau(k)$ denote by $m_\tau(\sigma)$ the number of $n-j-1$-simplices containing σ in X_τ. Note that $m_\tau(\sigma) = m(\tau\sigma)$ and by our previous assumption, $m_\tau(\sigma) \geq 1$ for every σ.
3. For a simplex $\sigma \in \Sigma_\tau(k)$, denote by $\Gamma_{\tau\sigma}$ the stabilizer of σ in Γ_τ.
4. For $0 \leq k \leq n-j-1$, denote by $C^k(X_\tau,\rho_\tau)$ the space of simplicial k-cochains of X_τ which are twisted by ρ_τ.

We also define $L^k(X_\tau,\rho_\tau), d_\tau, \delta_\tau, \Delta^+_\tau, \Delta^-_\tau, \Delta_\tau$ and the L^2-cohomology as before. Note that since X_τ is finite we get that $L^k(X_\tau,\rho_\tau) = C^k(X_\tau,\rho_\tau)$.

Define the *localization map*

$$C^k(X,\rho) \to C^{k-j-1}(X_\tau,\rho_\tau),\ \phi \to \phi_\tau$$

Where $\phi_\tau(\sigma) = \phi(\tau\sigma)$.

2.4 Further results from [BŚ97], [DJ00]

Proposition 2.3. *In [BŚ97],[DJ00] following results were proven:*

1. *[BŚ97, Lemma 1.3], [DJ00, Lemma 3.3] For $0 \le l < k \le n$, let $f = f(\tau,\sigma)$ be a Γ-invariant function on the set of pairs (τ,σ), where τ is an ordered l-simplex and σ is an ordered k-simplex with $\tau \subset \sigma$ Then*

$$\sum_{\sigma\in\Sigma(k,\Gamma)} \sum_{\substack{\tau\in\Sigma(l)\\ \tau\subset\sigma}} \frac{f(\tau,\sigma)}{|\Gamma_\sigma|} = \sum_{\tau\in\Sigma(l,\Gamma)} \sum_{\substack{\sigma\in\Sigma(k)\\ \tau\subset\sigma}} \frac{f(\tau,\sigma)}{|\Gamma_\tau|}$$

2. *[BŚ97, equality (1.9)] For $\psi \in C^l(X_\tau,\rho_\tau)$ we have*

$$\|\psi\|^2 = \sum_{\eta\in\Sigma_\tau(l,\Gamma_\tau)} \frac{m_\tau(\eta)}{|\Gamma_{\tau\eta}|\,(l+1)!}\,|\psi(\eta)|^2 = \frac{1}{(l+1)!\,|\Gamma_\tau|}\sum_{\eta\in\Sigma_\tau(l)} m(\tau\eta)\,|\psi(\eta)|^2$$

3. *[BŚ97, Lemma 1.11] For $0 \le k \le n$, $\tau \in \Sigma(k-1)$ and $\phi \in L^k(X,\rho)$, denote by ϕ^0_τ the component of ϕ_τ in the subspace of constant maps in $C^0(X_\tau,\rho_\tau)$ then*

$$\left\|\phi^0_\tau\right\|^2 = \frac{m(\tau)}{(n-k+1)\,|\Gamma_\tau|}\,|\delta\phi(\tau)|^2$$

and therefore

$$\|\delta\phi\|^2 = \sum_{\tau\in\Sigma(k-1,\Gamma)} \frac{n-k+1}{k!}\left\|\phi^0_\tau\right\|^2$$

4. *[BŚ97, Theorem 1.12] Let $0 \le j < k < n$ and $\phi \in L^k(X,\rho)$ then*

$$k!(\|d\phi\|^2 - (n-k)\|\phi\|^2) = (k-j-1)!\sum_{\tau\in\Sigma(j,\Gamma)} \|d_\tau\phi_\tau\|^2 - (n-k)\|\phi_\tau\|^2$$

[BŚ97, Corollary 1.13] For $0 < k < n$ and $\tau \in \Sigma(k-1)$ define a quadratic form Q_τ on $C^0(X_\tau,\rho_\tau)$ by

$$Q_\tau(\psi) = \|d_\tau\psi\|^2 - \frac{k}{k+1}(n-k)\,\|\psi\|^2 = \left\langle \Delta^+_\tau\psi,\psi\right\rangle - \frac{k}{k+1}(n-k)\,\|\psi\|^2$$

Then for every $\phi \in L^k(X,\rho)$ we have

$$k!\,\|d\phi\|^2 = \sum_{\tau\in\Sigma(k-1,\Gamma)} Q_\tau(\phi_\tau)$$

2.5 Vanishing of L^2-cohomologies

The idea (taken from [BŚ97]) for proving that $L^2 H^k(X,\rho) = 0$ for every ρ goes as follows: prove that there is an $\varepsilon > 0$ such that:

$$\forall \phi \in L^k(X,\rho), \|d\phi\|^2 = 0 \Rightarrow \|\delta\phi\|^2 \geq \varepsilon \|\phi\|^2$$

This will imply that $ker(d \mid L^k(X,\rho)) \cap ker(\delta \mid L^k(X,\rho)) = 0$ and therefore $ker(\Delta^+ \mid L^k(X,\rho)) \cap ker(\Delta^- \mid L^k(X,\rho)) = 0$ so we get $L^2\mathcal{H}^k(X,\rho) = 0$. Also, by the above inequality, we get that the image of $\Delta^- \mid L^k(X,\rho)$ is closed in $ker(\Delta^+ \mid L^k(X,\rho))$. Hence $ker(\Delta^+ \mid L^k(X,\rho)) = im(\Delta^- \mid L^k(X,\rho))$ and therefore $L^2 H^k(X,\rho) = 0$. In particular, one could prove the vanishing of $L^k(X,\rho)$ for every ρ by showing that there is an $\varepsilon > 0$ such that for every representation ρ the following holds:

$$\forall \phi \in L^k(X,\rho), \|d\phi\|^2 + \|\delta\phi\|^2 \geq \varepsilon \|\phi\|^2$$

In the same manner, to prove that $L^2\mathcal{H}^k(X,\rho) = 0$ for every ρ, it is enough to prove:

$$\forall \phi \in L^k(X,\rho), \|\delta\phi\|^2 = 0 \Rightarrow \|d\phi\|^2 > 0$$

3 Criteria via Laplacian eigenvalues

In this section we shall find criteria for the vanishing of L^2-cohomologies using the eigenvalues for the Laplacians.

Definition 3.1. *For $0 \leq k < l$ integers define M_k^l as following: let γ be an ordered simplex of dimension l denote $F = \{\sigma \sqsubset \gamma : dim(\sigma) = k\}, F' = \{\nu \sqsubset \gamma : dim(\nu) = k+1\}$. Denote by $[\nu : \sigma]$ the sign of σ in ν, i.e. if $\nu = (v_0, ..., v_{k+1}), \sigma = (v_0, ..., \hat{v_i}, ..., v_{k+1})$, then $[\nu : \sigma] = (-1)^i$. Now M_k^l is a matrix indexed by $F \cup F'$ defined as*

$$(M_k^l)_{\alpha,\beta} = \begin{cases} x_\alpha & \alpha = \beta \in F \\ [\beta : \alpha] & \alpha \in F, \beta \in F', \alpha \sqsubset \beta \\ [\alpha : \beta] & \beta \in F, \alpha \in F', \beta \sqsubset \alpha \\ 0 & otherwise \end{cases}$$

Define $p_k^l(x_\sigma)$ to be the multi polynomial with variables indexed by F as

$$p_k^l(x_\sigma) = det(M_k^l)$$

Example 3.2. *For $l = k+1$,*

$$p_k^l(x_\sigma) = \sum_{\sigma \in F} \prod_{\sigma' \in F, \sigma' \neq D} x_{\sigma'}$$

notice that in this case we get

$$p_k^l(x_\sigma) = \nabla \cdot (\prod_{\sigma' \in F} x_{\sigma'})$$

Remark 3.3. *Notice that if* $2(k+1) < l+1$ *then*

$$|F| = \left(\begin{array}{c} l+1 \\ k+1 \end{array} \right) \leq \left(\begin{array}{c} l+1 \\ k+2 \end{array} \right) = |F'|$$

and therefore p_k^l *is constant (if* $2(k+1) < l$ *then* $p_k^l \equiv 0$*).*

For every simplex τ of dimension $k-1$ denote by λ_τ the smallest positive eigenvalue of Δ_τ^+ on X_τ (the link of τ) also denote $\overline{\lambda_\tau} = \lambda_\tau - \frac{k(n-k)}{k+1}$. Now for a simplex σ of dimension k denote

$$S_\sigma = \sum_{\tau \in \Sigma(k-1), \tau \sqsubset \sigma} \overline{\lambda_\tau}$$

Theorem 3.4. *If there is an* $l > k$ *and* $\varepsilon > 0$ *such that for any* $\gamma \in \Sigma(l, \Gamma)$ *one has*

$$p_k^l(\lambda - S_\sigma) = 0 \Rightarrow \lambda \geq \varepsilon$$

where $p_k^l(\lambda - S_\sigma)$ *is the polynomial obtained by placing* $x_\sigma = \lambda - S_\sigma$*, then* $L^2 H^k(X, \rho) = 0$ *for every* ρ*.*

Proof. We shall start with repeating the proof in [BŚ97]:
Let $\phi \in L^k(X, \rho)$ then by Proposition 2.3 (4), one has

$$k! \left\| d\phi \right\|^2 = \sum_{\tau \in \Sigma(k-1, \Gamma)} \langle \Delta_\tau^+(\phi_\tau), \phi_\tau \rangle - \frac{k}{k+1}(n-k) \left\| \phi_\tau \right\|^2$$

Now denote as before ϕ_τ^0 to be the projection of ϕ_τ on the space of constant function on X_τ and by ϕ_τ^1 its orthogonal complement. Note that since $ker(\Delta_\tau^+)$ is that space of constant function on X_τ (since X_τ is connected), we get that

$$\langle \Delta_\tau^+(\phi_\tau), \phi_\tau \rangle = \langle \Delta_\tau^+(\phi_\tau^1), \phi_\tau^1 \rangle \geq \lambda_\tau \|\phi_\tau^1\|^2$$

Therefore

$$k! \left\| d\phi \right\|^2 \geq \sum_{\tau \in \Sigma(k-1, \Gamma)} \lambda_\tau \|\phi_\tau^1\|^2 - \frac{k}{k+1}(n-k) \left\| \phi_\tau \right\|^2 =$$

$$= \sum_{\tau \in \Sigma(k-1, \Gamma)} \lambda_\tau \|\phi_\tau\|^2 - \lambda_\tau \|\phi_\tau^0\|^2 - \frac{k}{k+1}(n-k) \left\| \phi_\tau \right\|^2$$

So we get

$$k! \left\| d\phi \right\|^2 + \sum_{\tau \in \Sigma(k-1, \Gamma)} \lambda_\tau \|\phi_\tau^0\|^2 \geq \sum_{\tau \in \Sigma(k-1, \Gamma)} \overline{\lambda_\tau} \left\| \phi_\tau \right\|^2$$

Note that since $\|d_\tau\|^2 \leq 2(n-k)$ for every τ (see [BŚ97, Proposition 1.5]) then $\lambda_\tau \leq 2(n-k)$ for every τ and therefore

$$k! \frac{2(n-k)}{n-k+1} \|\delta\phi\|^2 = \sum_{\tau \in \Sigma(k-1, \Gamma)} 2(n-k) \|\phi_\tau^0\|^2 \geq \sum_{\tau \in \Sigma(k-1, \Gamma)} \lambda_\tau \|\phi_\tau^0\|^2$$

So we get

$$k!\left\|d\phi\right\|^2+k!\frac{2(n-k)}{n-k+1}\|\delta\phi\|^2\geq\sum_{\tau\in\Sigma(k-1,\Gamma)}\overline{\lambda_\tau}\left\|\phi_\tau\right\|^2$$

Remark 3.5. *At this stage, if we wanted to prove the result in [BŚ97] (every $\overline{\lambda_\tau}\geq\varepsilon$ implies the vanishing of the cohomology) we would be done, because then*

$$k!\left\|d\phi\right\|^2+k!\frac{2(n-k)}{n-k+1}\|\delta\phi\|^2\geq\varepsilon(k+1)!\|\phi\|^2$$

Now we shall assume that $\phi\in ker(d)$ and show that under the condition stated in the theorem we get that

$$\frac{2(n-k)}{n-k+1}\|\delta\phi\|^2\geq\varepsilon\|\phi\|^2$$

and that will finish the proof at stated at the beginning of this section.

By definition and Proposition 2.3 (2) we get that

$$\sum_{\tau\in\Sigma(k-1,\Gamma)}\overline{\lambda_\tau}\left\|\phi_\tau\right\|^2=\sum_{\tau\in\Sigma(k-1,\Gamma)}\overline{\lambda_\tau}\frac{1}{|\Gamma_\tau|}\sum_{\eta\in\Sigma_\tau(0)}m(\tau\eta)\left|\phi(\tau\eta)\right|^2=$$

$$=\sum_{\tau\in\Sigma(k-1,\Gamma)}\overline{\lambda_\tau}\frac{1}{|\Gamma_\tau|}\sum_{\sigma\in\Sigma(k),\tau\subset\sigma}\frac{1}{(k+1)!}m(\sigma)\left|\phi(\sigma)\right|^2$$

by Proposition 2.3 (1) we can change the order of summation and get

$$\sum_{\sigma\in\Sigma(k,\Gamma)}\frac{m(\sigma)|\phi(\sigma)|^2}{(k+1)!|\Gamma_\sigma|}\sum_{\tau\in\Sigma(k-1),\tau\subset\sigma}\overline{\lambda_\tau}=k!\sum_{\sigma\in\Sigma(k,\Gamma)}\frac{m(\sigma)|\phi(\sigma)|^2}{(k+1)!|\Gamma_\sigma|}S_\sigma$$

Now note that

$$(l+1)!\begin{pmatrix}n-k\\l-k\end{pmatrix}m(\sigma)=\sum_{\gamma\in\Sigma(l),\sigma\subset\gamma}m(\gamma)$$

so we get

$$k!\sum_{\sigma\in\Sigma(k,\Gamma)}\frac{m(\sigma)|\phi(\sigma)|^2}{(k+1)!|\Gamma_\sigma|}S_\sigma=k!\sum_{\sigma\in\Sigma(k,\Gamma)}\frac{|\phi(\sigma)|^2}{(k+1)!|\Gamma_\sigma|}S_\sigma\sum_{\gamma\in\Sigma(l),\sigma\subset\gamma}\frac{m(\gamma)}{(l+1)!\begin{pmatrix}n-k\\l-k\end{pmatrix}}$$

and by changing the order of summation again we get

$$\frac{k!}{(k+1)!}\sum_{\gamma\in\Sigma(l,\Gamma)}\frac{m(\gamma)}{(l+1)!\begin{pmatrix}n-k\\l-k\end{pmatrix}|\Gamma_\gamma|}\sum_{\sigma\in\Sigma(k),\sigma\subset\gamma}S_\sigma|\phi(\sigma)|^2$$

Now we shall show that, under the conditions of the theorem, for every $\gamma\in\Sigma(l,\Gamma)$ we have the the following inequality

$$\sum_{\sigma\in\Sigma(k),\sigma\subset\gamma}S_\sigma|\phi(\sigma)|^2\geq\varepsilon\sum_{\sigma\in\Sigma(k),\sigma\subset\gamma}|\phi(\sigma)|^2$$

and that will complete the proof because then we have

$$\frac{k!}{(k+1)!}\sum_{\gamma\in\Sigma(l,\Gamma)}\frac{m(\gamma)}{(l+1)!\begin{pmatrix} n-k \\ l-k \end{pmatrix}|\Gamma_\gamma|}\sum_{\sigma\in\Sigma(k),\sigma\subset\gamma}S_\sigma|\phi(\sigma)|^2\geq$$

$$\geq\frac{k!}{(k+1)!}\sum_{\gamma\in\Sigma(l,\Gamma)}\frac{m(\gamma)}{(l+1)!\begin{pmatrix} n-k \\ l-k \end{pmatrix}|\Gamma_\gamma|}\sum_{\sigma\in\Sigma(k),\sigma\subset\gamma}\varepsilon|\phi(\sigma)|^2=$$

$$=\varepsilon\frac{k!}{(k+1)!}\sum_{\sigma\in\Sigma(k,\Gamma)}\frac{|\phi(\sigma)|^2}{|\Gamma_\sigma|}\sum_{\gamma\in\Sigma(l),\sigma\subset\gamma}\frac{m(\gamma)}{(l+1)!\begin{pmatrix} n-k \\ l-k \end{pmatrix}}=$$

$$=\varepsilon k!\sum_{\sigma\in\Sigma(k,\Gamma)}\frac{m(\sigma)|\phi(\sigma)|^2}{(k+1)!|\Gamma_\sigma|}=\varepsilon k!\|\phi\|^2$$

And therefore we get

$$\frac{2(n-k)}{n-k+1}\|\delta\phi\|^2\geq\varepsilon\|\phi\|^2$$

So we are left with proving the following inequality - for every $\phi\in L^k(X,\rho)\cap ker(d)$ and for every $\gamma\in\Sigma(l,\Gamma)$ one has (under the conditions of the theorem):

$$\sum_{\sigma\in\Sigma(k),\sigma\subset\gamma}S_\sigma|\phi(\sigma)|^2\geq\varepsilon\sum_{\sigma\in\Sigma(k),\sigma\subset\gamma}|\phi(\sigma)|^2$$

Fix $\gamma\in\Sigma(l,\Gamma)$, first note that since ϕ is alternating we get that

$$\sum_{\sigma\in\Sigma(k),\sigma\subset\gamma}S_\sigma|\phi(\sigma)|^2=(k+1)!\sum_{\sigma\in\Sigma(k),\sigma\sqsubset\gamma}S_\sigma|\phi(\sigma)|^2$$

and

$$\sum_{\sigma\in\Sigma(k),\sigma\subset\gamma}|\phi(\sigma)|^2=(k+1)!\sum_{\sigma\in\Sigma(k),\sigma\sqsubset\gamma}|\phi(\sigma)|^2$$

therefore it is enough to prove that

$$\sum_{\sigma\in\Sigma(k),\sigma\sqsubset\gamma}S_\sigma|\phi(\sigma)|^2\geq\varepsilon\sum_{\sigma\in\Sigma(k),\sigma\sqsubset\gamma}|\phi(\sigma)|^2$$

Now we shall need the following simple but useful lemma, which is a straightforward generalization of [BŚ97, Lemma 2.3].

Lemma 3.6. *For a finite set $F=\{x_1,...,x_n\}$, denote by $C(F,H)$ the space of maps from F to H. On that space there is a natural inner product:*

$$\langle\phi,\psi\rangle=\sum_{k=1}^{n}\langle\phi(x_k),\psi(x_k)\rangle$$

and denote by $\|.\|$ the norm induced by this inner product. Note that $C(F,H) = C(F,\mathbb{R}) \otimes H$ (as a Hilbert space). Let $T_j, j = 1, ..., m$ and $M_i, i = 1, ..., l$ be bounded linear operators on $C(F,H)$ so there are operators $T_{j,\mathbb{R}}, M_{i,\mathbb{R}}$ on $C(F,\mathbb{R})$ such that $T_j = T_{j,\mathbb{R}} \otimes id, M_i = M_{i,\mathbb{R}} \otimes id$. Let $a_1, ..., a_j \in \mathbb{R}$ be constants, then if there is an ε such that

$$\forall \phi \in C(F,\mathbb{R}), \phi \in \cap ker(M_{i,\mathbb{R}}) \Rightarrow \sum_j a_j \|T_{j,\mathbb{R}} \phi\|^2 \geq \varepsilon \|\phi\|^2$$

Then also

$$\forall \phi \in C(F,H), \phi \in \cap ker(M_i) \Rightarrow \sum_j a_j \|T_j \phi\|^2 \geq \varepsilon \|\phi\|^2$$

Proof. Choose $\{v_\alpha\}$ orthonormal basis of H. For every $\phi \in C(F,H)$ one has $\phi_\alpha \in C(X,\mathbb{R})$ such that $\phi = \sum \phi_\alpha \otimes v_\alpha$. Now note that $\phi \in \cap ker(M_i)$ iff $\forall \alpha, \phi_\alpha \in \cap ker(M_{i,\mathbb{R}})$, so we get

$$\forall \phi \in \cap ker(M_i), \sum_j a_j \|T_j \phi\|^2 = \| \sum_j T_j \sum_\alpha \phi_\alpha \otimes v_\alpha \|^2 = \sum_\alpha \sum_j a_j \|T_{j,\mathbb{R}} \phi_\alpha\|^2 \geq$$

$$\geq \sum_\alpha \varepsilon \|\phi_\alpha\|^2 = \varepsilon \|\phi\|^2$$

□

Now we can use the above lemma to reduce our problem to a much simpler one. Denote $F = \{\sigma \in \Sigma(k) : \sigma \sqsubset \gamma\}$ and $F' = \{\nu \in \Sigma(k+1) : \nu \sqsubset \gamma\}$ and for $\phi \in L^k(X,\rho) \cap ker(d)$ we can look at the restriction of ϕ to F. Note that $\phi \in ker(d)$ implies that for every $\nu \in \Sigma(k+1)$ we have $d\phi(\nu) = 0$ and in particular $\forall \nu \in F', d\phi(\nu) = 0$. Therefore for every $\nu \in F'$, we define M_ν acting on $C(F,H)$ as $M_\nu \phi = \sum (-1)^i \phi(\nu_i)$. For every $\sigma \in F$ define T_σ acting on $C(F,H)$ as the projection on the space spanned by the indicator function of σ. If we can prove that for every $\phi \in C(F,H)$ the following holds

$$\phi \in \cap ker(M_\nu) \Rightarrow \sum_{\sigma \sqsubset \gamma} S_\sigma \|T_\sigma \phi\|^2 \geq \varepsilon \|\phi\|^2$$

then in particular, for every $\phi \in C^k(X,\rho)$ we have

$$\sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} S_\sigma |\phi(\sigma)|^2 \geq \varepsilon \sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} |\phi(\sigma)|^2$$

By the above lemma, it is enough to prove that for $\phi \in C(F,\mathbb{R})$ we have

$$\phi \in \cap ker(M_\nu) \Rightarrow \sum_{\sigma \sqsubset \gamma} S_\sigma \|T_\sigma \phi\|^2 \geq \varepsilon \|\phi\|^2$$

and if we denote $\phi(\sigma) = x_\sigma$ we get the following problem: prove that

$$\sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} S_\sigma x_\sigma^2 \geq \varepsilon \sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} x_\sigma^2$$

under the constraints

$$\forall \nu \in F', \sum_{i=0}^{k+1} (-1)^i x_{\nu_i} = 0$$

since both sides of the inequality we are trying to prove are quadratic, WLOG it is enough to prove that

$$\sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} S_\sigma x_\sigma^2 \geq \varepsilon$$

under the constraints

$$\sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} x_\sigma^2 = 1$$

$$\forall \nu \in F', \sum_{\sigma \in F, \sigma \sqsubset \nu} [\nu : \sigma] x_\sigma = 0$$

This is a problem of finding a minimum of a function in $\mathbb{R}^{|F|}$ under constraints which define a compact set in $\mathbb{R}^{|F|}$ and so we can use the Lagrange multiplier theorem. Define the Lagrange function to be

$$\Lambda(x_\sigma, \mu_\nu, \lambda) = \sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} S_\sigma x_\sigma^2 - 2 \sum_{\nu \in F'} \mu_\nu (\sum_{\sigma \in F, \sigma \sqsubset \nu} [\nu : \sigma] x_\sigma - \lambda (\sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} x_\sigma^2 - 1)$$

(the 2 multiplying $\sum_{\nu \in F'}$ is added for convenience)

So for every $\sigma \in F$ we get a equation by derivation of Λ by x_σ:

$$2 S_\sigma x_\sigma - 2\lambda x_\sigma - 2 \sum_{\nu \in F', \sigma \sqsubset \nu} [\nu : \sigma] \mu_\nu = 0$$

if we multiply every such equation by x_σ and add them up (over all $\sigma \in F$) we get

$$2 \sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} S_\sigma x_\sigma^2 - 2\lambda (\sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} x_\sigma^2) - 2 \sum_{\nu \in F'} \mu_\nu \sum_{\sigma \in F, \sigma \sqsubset \nu} [\nu : \sigma] x_\sigma = 0$$

and by the equations coming from the constraints we get

$$\sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} S_\sigma x_\sigma^2 = \lambda$$

So the minimum is must some λ which is a part of a vector $(x_\sigma, \mu_\nu, \lambda)$ which solves $\nabla \Lambda = 0$.

Treat λ as a parameter and consider the system of linear equations in (x_σ, μ_ν)

$$(\lambda - S_\sigma) x_\sigma + \sum_{\nu \in F', \sigma \sqsubset \nu} [\nu : \sigma] \mu_\nu = 0$$

$$\sum_{\sigma \in F, \sigma \sqsubset \nu} [\nu : \sigma] x_\sigma = 0$$

Note that from

$$\sum_{\sigma \in \Sigma(k), \sigma \sqsubset \gamma} x_\sigma^2 = 1$$

the minimum is obtain only if this system of equations have a non trivial solution, that is only if the determinant is zero, but this determinant is exactly $p_k^l(\lambda - S_\sigma)$ and by the conditions of the theorem we know that for every root λ of this polynomial we have $\lambda \geq \varepsilon$ and so we are done.

□

Now Theorem 1 is proven as a corollary:

Corollary 3.7. *Let Γ be a locally compact, properly discontinuous, unimodular group of automorphisms of X acting cocompactly on X such that X is a locally finite contractible 2-dimensional simplicial complex. If for every $(u, v, w) \in \Sigma(2, \Gamma)$ the following holds:*

- $\overline{\lambda_u} + \overline{\lambda_v} + \overline{\lambda_w} > 0$
- $S_{(u,v)}S_{(u,w)} + S_{(u,v)}S_{(v,w)} + S_{(u,w)}S_{(v,w)} > 0$

then Γ has property (T).

Proof. By the above theorem it is enough to prove that for every $(u, v, w) \in \Sigma(2, \Gamma)$ all the roots of $p_1^2(\lambda - S_{(u,v)}, \lambda - S_{(v,w)}, \lambda - S_{(u,w)})$ are positive. We have

$$p_1^2(\lambda - S_{(u,v)}, \lambda - S_{(v,w)}, \lambda - S_{(u,w)}) =$$

$$= (\lambda - S_{(u,v)})(\lambda - S_{(v,w)}) + (\lambda - S_{(u,v)})(\lambda - S_{(u,w)}) + (\lambda - S_{(v,w)})(\lambda - S_{(u,w)})$$

one can find the roots of the polynomial explicitly and the roots of this polynomial are positive iff

- $\overline{\lambda_u} + \overline{\lambda_v} + \overline{\lambda_w} > 0$
- $S_{(u,v)}S_{(u,w)} + S_{(u,v)}S_{(v,w)} + S_{(u,w)}S_{(v,w)} > 0$

□

4 Criteria via cosine of links

We begin with a general definition for a cosine of a finite simplicial complex with respect to a group of automorphisms.

Definition 4.1. *Let Y be a connected finite simplicial complex, Λ be a subgroup of $Aut(Y)$ and ρ be a unitary representation of Λ. Define the cosine of Y with respect to Λ and ρ to be*

$$cos(Y, \Lambda, \rho) = sup\{\frac{\sum_{(u,v) \in \Sigma(1,\Lambda)} m((u,v)) \langle \phi^1(u), \phi^1(v) \rangle}{\sum_{(u,v) \in \Sigma(1,\Lambda)} m((u,v)) |\phi(u)||\phi(v)|} : 0 \neq \phi \in C^0(Y, \rho)\}$$

where ϕ^1 is (as above) the projection of ϕ on the orthogonal compliment of the space of constant functions. Define also

$$cos(Y,\Lambda) = sup\{cos(Y,\Lambda,\rho) : \rho \text{ is a unitary representation}\}$$

Remark 4.2. *Note that $\Lambda_1 < \Lambda_2$ implies that $cos(Y,\Lambda_1) \geq cos(Y,\Lambda_2)$ and in particular if we denote $cos(Y) = cos(Y,\{e\})$ then $cos(Y) \geq cos(Y,\Lambda)$ for every Λ.*

Now we return to our general setting, i.e. X is a simplicial complex, Γ is a locally compact unimodular group of automorphisms of X acting properly discontinuously on X and so on.

Definition 4.3. *For $\gamma \in \Sigma(k+1)$ define $A(\gamma,\Gamma,\rho)$ to be a matrix indexed by $\{\sigma \in \Sigma(k) : \sigma \sqsubset \gamma\}$ as*

$$(A(\gamma,\Gamma,\rho))_{\alpha,\beta} = \begin{cases} 1 \,;\ \alpha = \beta \\ -cos(X_{\alpha\cap\beta},\Gamma_{\alpha\cap\beta},\rho) \,;\ \alpha \neq \beta \end{cases}$$

where $\alpha \cap \beta$ is the simplex of dimension $k-1$ that is contained in α and in β. In the same way, define

$$(A(\gamma,\Gamma))_{\alpha,\beta} = \begin{cases} 1 \,;\ \alpha = \beta \\ -cos(X_{\alpha\cap\beta},\Gamma_{\alpha\cap\beta}) \,;\ \alpha \neq \beta \end{cases}$$

Denote $A(\gamma) = A(\gamma,\{e\})$.

Theorem 4.4. *If for every unitary representation ρ there is an $\varepsilon > 0$ such that for every $\gamma \in \Sigma(k+1,\Gamma)$ the matrix $A(\gamma,\Gamma,\rho)$ is positive definite with eigenvalues greater or equal to ε then $L^2H^k(X,\rho) = 0$ for every ρ.*

Proof. Fix ρ and $\phi \in L^k(X,\rho)$. For every $\gamma \in \Sigma(k+1,\Gamma)$ define the vector $V_\gamma = (|\phi(\sigma)| : \sigma \in \Sigma(k), \sigma \sqsubset \gamma)$ so by the conditions of the theorem we get

$$V_\gamma^t A(\gamma,\Gamma,\rho) V_\gamma \geq \varepsilon \sum_{\sigma\in\Sigma(k),\sigma\sqsubset\gamma} |\phi(\sigma)|^2$$

which implies

$$\sum_{\gamma\in\Sigma(k+1,\Gamma)} \frac{m(\gamma)}{(k+2)!|\Gamma_\gamma|} V_\gamma^t A(\gamma,\Gamma,\rho) V_\gamma \geq \varepsilon \sum_{\gamma\in\Sigma(k+1,\Gamma)} \frac{m(\gamma)}{|\Gamma_\gamma|(k+2)!(k+1)!} \sum_{\sigma\in\Sigma(k),\sigma\subset\gamma} |\phi(\sigma)|^2$$

changing the order of summation in the right hand side of the inequality yields

$$\varepsilon \sum_{\sigma\in\Sigma(k,\Gamma)} \frac{|\phi(\sigma)|^2}{|\Gamma_\sigma|(k+2)!(k+1)!} \sum_{\gamma\in\Sigma(k+1),\sigma\subset\gamma} m(\gamma) =$$

$$= \varepsilon(n-k) \sum_{\sigma\in\Sigma(k,\Gamma)} \frac{m(\sigma)|\phi(\sigma)|^2}{|\Gamma_\sigma|(k+1)!} = \varepsilon(n-k)\|\phi\|^2$$

Now we shall work on the left hand side of the above inequality, i.e. on

$$\sum_{\gamma\in\Sigma(k+1,\Gamma)} \frac{m(\gamma)}{(k+2)!|\Gamma_\gamma|} V_\gamma^t A(\gamma,\Gamma,\rho)V_\gamma$$

writing the term $V_\gamma^t A(\gamma,\Gamma,\rho)V_\gamma$ explicitly gives

$$\sum_{\gamma\in\Sigma(k+1,\Gamma)} \frac{m(\gamma)}{(k+2)!|\Gamma_\gamma|} \sum_{\sigma\in\Sigma(k),\sigma\sqsubset\gamma} |\phi(\sigma)|^2 -$$

$$-\sum_{\gamma\in\Sigma(k+1,\Gamma)} \frac{m(\gamma)}{(k+2)!|\Gamma_\gamma|} \sum_{\substack{\sigma,\sigma'\in\Sigma(k)\\ \sigma,\sigma'\sqsubset\gamma,\sigma\neq\sigma'}} cos(X_{\sigma\cap\sigma'},\Gamma_{\sigma\cap\sigma'},\rho)|\phi(\sigma)||\phi(\sigma')|$$

First we look at

$$\sum_{\gamma\in\Sigma(k+1,\Gamma)} \frac{m(\gamma)}{(k+2)!|\Gamma_\gamma|} \sum_{\sigma\in\Sigma(k),\sigma\sqsubset\gamma} |\phi(\sigma)|^2 =$$

$$= \sum_{\gamma\in\Sigma(k+1,\Gamma)} \frac{m(\gamma)}{(k+2)!(k+1)!|\Gamma_\gamma|} \sum_{\sigma\in\Sigma(k),\sigma\subset\gamma} |\phi(\sigma)|^2 =$$

$$= \sum_{\sigma\in\Sigma(k,\Gamma)} \frac{|\phi(\sigma)|^2}{(k+2)!(k+1)!|\Gamma_\sigma|} \sum_{\gamma\in\Sigma(k+1),\sigma\subset\gamma} m(\gamma) =$$

$$(n-k)(k+2)! \sum_{\sigma\in\Sigma(k,\Gamma)} \frac{m(\sigma)|\phi(\sigma)|^2}{(k+2)!(k+1)!|\Gamma_\sigma|} = (n-k)\|\phi\|^2$$

Now we shall look at

$$\sum_{\gamma\in\Sigma(k+1,\Gamma)} \frac{m(\gamma)}{(k+2)!|\Gamma_\gamma|} \sum_{\substack{\sigma,\sigma'\in\Sigma(k)\\ \sigma,\sigma'\sqsubset\gamma,\sigma\neq\sigma'}} cos(X_{\sigma\cap\sigma'},\Gamma_{\sigma\cap\sigma'},\rho)|\phi(\sigma)||\phi(\sigma')| =$$

$$= \sum_{\gamma\in\Sigma(k+1,\Gamma)} \frac{m(\gamma)}{(k+2)!|\Gamma_\gamma|} \sum_{\tau\in\Sigma(k-1),\tau\sqsubset\gamma} cos(X_\tau,\Gamma_\tau,\rho) \sum_{\substack{u,v\in\Sigma(0)\\ \tau u,\tau v\subset\gamma}} |\phi(\tau u)||\phi(\tau v)| =$$

$$= \frac{1}{k!} \sum_{\gamma\in\Sigma(k+1,\Gamma)} \frac{m(\gamma)}{(k+2)!|\Gamma_\gamma|} \sum_{\tau\in\Sigma(k-1),\tau\subset\gamma} cos(X_\tau,\Gamma_\tau,\rho) \sum_{\substack{u,v\in\Sigma(0)\\ \tau u,\tau v\subset\gamma}} |\phi(\tau u)||\phi(\tau v)| =$$

$$= \frac{1}{k!} \sum_{\tau\in\Sigma(k-1,\Gamma)} \frac{cos(X_\tau,\Gamma_\tau,\rho)}{(k+2)!|\Gamma_\tau|} \sum_{\gamma\in\Sigma(k+1),\tau\subset\gamma} m(\gamma) \sum_{\substack{u,v\in\Sigma(0)\\ \tau u,\tau v\subset\gamma}} |\phi(\tau u)||\phi(\tau v)| =$$

$$= \frac{1}{k!} \sum_{\tau\in\Sigma(k-1,\Gamma)} \frac{cos(X_\tau,\Gamma_\tau,\rho)}{|\Gamma_\tau|} \sum_{(u,v)\in\Sigma_\tau(1)} m_\tau((u,v))|\phi_\tau(u)||\phi_\tau(v)|$$

By the definition of $cos(X_\tau, \Gamma_\tau, \rho)$ we get

$$\frac{1}{k!} \sum_{\tau \in \Sigma(k-1,\Gamma)} \frac{cos(X_\tau, \Gamma_\tau, \rho)}{|\Gamma_\tau|} \sum_{(u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) |\phi_\tau(u)| |\phi_\tau(v)| \geq$$

$$\geq \frac{1}{k!} \sum_{\tau \in \Sigma(k-1,\Gamma)} \frac{1}{|\Gamma_\tau|} \sum_{(u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) \langle \phi_\tau^1(u), \phi_\tau^1(v) \rangle$$

Now note that

$$\sum_{(u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) \phi_\tau^1(u) = \sum_{u \in \Sigma_\tau(0)} \phi_\tau^1(u) \sum_{v \in \Sigma_\tau(0), (u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) =$$

$$= (n-k) \sum_{u \in \Sigma_\tau(0)} m_\tau(u) \phi_\tau^1(u) = 0$$

and

$$\frac{1}{|\Gamma_\tau|} \sum_{(u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) |\phi_\tau^0|^2 = \frac{1}{|\Gamma_\tau|} \sum_{u \in \Sigma_\tau(0)} |\phi_\tau^0|^2 \sum_{v \in \Sigma_\tau(0), (u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) =$$

$$= \frac{n-k}{|\Gamma_\tau|} \sum_{u \in \Sigma_\tau(0)} m_\tau(u) |\phi_\tau^0|^2 = (n-k) \|\phi_\tau^0\|^2$$

and therefore

$$\frac{1}{|\Gamma_\tau|} \sum_{(u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) \langle \phi_\tau(u), \phi_\tau(v) \rangle =$$

$$= \frac{1}{|\Gamma_\tau|} \sum_{(u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) \langle \phi_\tau^0 + \phi_\tau^1(u), \phi_\tau^0 + \phi_\tau^1(v) \rangle =$$

$$= (n-k) \|\phi_\tau^0\|^2 + \frac{1}{|\Gamma_\tau|} \sum_{(u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) \langle \phi_\tau^1(u), \phi_\tau^1(v) \rangle$$

so we get

$$\frac{1}{k!} \sum_{\tau \in \Sigma(k-1,\Gamma)} \frac{1}{|\Gamma_\tau|} \sum_{(u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) \langle \phi_\tau^1(u), \phi_\tau^1(v) \rangle =$$

$$= \frac{1}{k!} \sum_{\tau \in \Sigma(k-1,\Gamma)} -(n-k) \|\phi_\tau^0\|^2 + \frac{1}{|\Gamma_\tau|} \sum_{(u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) \langle \phi_\tau(u), \phi_\tau(v) \rangle =$$

$$= -\frac{n-k}{n-k+1} \|\delta\phi\|^2 + \frac{1}{k!} \sum_{\tau \in \Sigma(k-1,\Gamma)} \frac{1}{|\Gamma_\tau|} \sum_{(u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) \langle \phi_\tau(u), \phi_\tau(v) \rangle$$

Now we shall look at

$$\frac{1}{k!} \sum_{\tau \in \Sigma(k-1,\Gamma)} \frac{1}{|\Gamma_\tau|} \sum_{(u,v) \in \Sigma_\tau(1)} m_\tau((u,v)) \langle \phi_\tau(u), \phi_\tau(v) \rangle =$$

$$\frac{1}{k!}\sum_{\tau\in\Sigma(k-1,\Gamma)}\frac{1}{|\Gamma_\tau|}\sum_{(u,v)\in\Sigma_\tau(1)}[m_\tau((u,v))(-\frac{1}{2}|d_\tau\phi_\tau((u,v))|^2)+$$

$$+\frac{1}{2}m_\tau((u,v))|\phi(u)|^2+\frac{1}{2}m_\tau((u,v))|\phi(u)|^2]=$$

$$=\frac{1}{k!}\sum_{\tau\in\Sigma(k-1,\Gamma)}-\|d_\tau\phi_\tau\|^2+\frac{1}{k!}\sum_{\tau\in\Sigma(k-1,\Gamma)}\frac{1}{|\Gamma_\tau|}\sum_{(u,v)\in\Sigma_\tau(1)}m_\tau((u,v))|\phi(u)|^2=$$

$$=\frac{1}{k!}\sum_{\tau\in\Sigma(k-1,\Gamma)}-\|d_\tau\phi_\tau\|^2+(n-k)\|\phi_\tau\|^2=-\|d\phi\|^2+(n-k)\|\phi\|^2$$

where the last equality is by Proposition 2.3 (4). To sum up, we got that

$$\sum_{\gamma\in\Sigma(k+1,\Gamma)}\frac{m(\gamma)}{(k+2)!|\Gamma_\gamma|}\sum_{\substack{\sigma,\sigma'\in\Sigma(k)\\ \sigma,\sigma'\sqsubset\gamma,\sigma\neq\sigma'}}cos(X_{\sigma\cap\sigma'},\Gamma_{\sigma\cap\sigma'},\rho)|\phi(\sigma)||\phi(\sigma')|\geq$$

$$\geq-\frac{n-k}{n-k+1}\|\delta\phi\|^2-\|d\phi\|^2+(n-k)\|\phi\|^2$$

and therefore

$$\sum_{\gamma\in\Sigma(k+1,\Gamma)}\frac{m(\gamma)}{(k+2)!|\Gamma_\gamma|}\sum_{\sigma\in\Sigma(k),\sigma\sqsubset\gamma}|\phi(\sigma)|^2-$$

$$-\sum_{\gamma\in\Sigma(k+1,\Gamma)}\frac{m(\gamma)}{(k+2)!|\Gamma_\gamma|}\sum_{\substack{\sigma,\sigma'\in\Sigma(k)\\ \sigma,\sigma'\sqsubset\gamma,\sigma\neq\sigma'}}cos(X_{\sigma\cap\sigma'},\Gamma_{\sigma\cap\sigma'},\rho)|\phi(\sigma)||\phi(\sigma')|\leq$$

$$\leq(n-k)\|\phi\|^2+\frac{n-k}{n-k+1}\|\delta\phi\|^2+\|d\phi\|^2-(n-k)\|\phi\|^2=$$

$$\frac{n-k}{n-k+1}\|\delta\phi\|^2+\|d\phi\|^2$$

so we got that under the conditions of the theorem

$$\frac{n-k}{n-k+1}\|\delta\phi\|^2+\|d\phi\|^2\geq\varepsilon(n-k)\|\phi\|^2$$

and we are done.

□

Corollary 4.5. *If there is an $\varepsilon>0$ such that for every $\gamma\in\Sigma(k+1,\Gamma)$ the matrix $A(\gamma,\Gamma)$ is positive definite with eigenvalues greater or equal to ε then $L^2H^k(X,\rho)=0$ for every ρ. And in the same way, if there is an $\varepsilon>0$ such that for every $\gamma\in\Sigma(k+1,\Gamma)$ the matrix $A(\gamma)$ is positive definite with eigenvalues greater or equal to ε then $L^2H^k(X,\rho)=0$*

Corollary 4.6. *Define the reduced cosine as follows:*

$$cos_r(X,\Gamma,\rho) = sup\{\frac{\sum_{(u,v)\in\Sigma(1,\Gamma)} m((u,v))\langle\phi(u),\phi(v)\rangle}{\sum_{(u,v)\in\Sigma(1,\Gamma)} m((u,v))|\phi(u)||\phi(v)|} : 0 \neq \phi \in C^0(X,\Gamma), \phi = \phi^1\}$$

and let $A_r(\gamma,\Gamma)$ be the corresponding matrix. If for every $\gamma \in \Sigma(k+1,\Gamma)$ the matrix $A_r(\gamma,\Gamma)$ is positive definite then $L^2\mathcal{H}^k(X,\rho) = 0$ for every ρ.

Proof. We just repeat the above proof of the theorem with $0 \neq \phi \in ker(\delta)$ (which means $\phi = \phi^1$) and get $\|d\phi\|^2 > 0$. So $ker(\delta) \cap ker(d) = 0$ and we are done. □

Applying the above corollary to the 2-dimensional case gives Theorem 2.

Remark 4.7. *The results stated above where inspired by [Kas11], we do not reproduce the results of [Kas11] for dimension n larger than 2. In [Kas11] the cosine matrix refers always to the angles between the $n-1$ simplices to get property (T), but in our result the cosine matrix of those angles gives only the vanishing of the $n-1$ reduced cohomology and to get property (T) one needs to look at the cosine matrix of angles between 1 simplices.*

5 Examples

5.1 GABs

A generalized m-gon (or in another name, a 1-dimensional spherical building) is a connected bipartite graph $L = (V,E)$ of diameter m and girth $2m$ in which each vertex lies on at least two edges. Denote $V = V_1 \cup V_2$ were V_1, V_2 are the two sides of the graph (there is an edge between two vertices only if one belong to V_1 and the other to V_2). A generalized m-gon is said to have parameters (s,t) if every vertex in V_1 has valency $s+1$ and every vertex in V_2 has valency $t+1$. A generalized m-gon is called thick if $s \geq 2, t \geq 2$. A theorem by Feit and Higman [FH64] states that a thick m-gon exists only if $m = 2,3,4,6,8$. Moreover, Feit and Higman computed the smallest positive eigenvalue for the Laplacian on general m-gon of type (s,t) and those are given in the list below

1. For $m = 2$ the generalized m-gon is a complete bipartite graph and the smallest positive eigenvalue of the Laplacian is always 1 (is does not depend on (s,t)).
2. For $m = 3$ one always have $s = t$ and the generalized m-gon is the flag complex of a projective plane. In that case the smallest positive eigenvalue of the Laplacian is
$$1 - \frac{\sqrt{s}}{s+1}$$
3. For $m = 4$ the smallest positive eigenvalue of the Laplacian is
$$1 - \sqrt{\frac{s+t}{(s+1)(t+1)}}$$

4. For $m = 6$ the smallest positive eigenvalue of the Laplacian is

$$1 - \sqrt{\frac{s + t + \sqrt{st}}{(s + 1)(t + 1)}}$$

5. For $m = 8$ the smallest positive eigenvalue of the Laplacian is

$$1 - \sqrt{\frac{s + t + \sqrt{2st}}{(s + 1)(t + 1)}}$$

"Geometries that are almost buildings" (GABs) and "Chamber system that are almost buildings" (SCABs) were introduced by Tits in [Tit81] (other names for those are "geometries of type M" and "chamber systems of type M"). GAB and SCABs can be viewed as finite n-dimensional simplicial complexes whose links of codimension 2 are generalized polygons (exact definition of GABs and SCABs can be found in [Ros86] and in [Tit81]). Below we introduce three examples of GABs and SCABs given in [Kan81] and in [Ros86], whose universal cover are an exotic affine buildings (i.e. affine building that do not arise from a local field) and whose fundamental groups acts on those buildings. These examples are interesting in our context because they fail to meet Żuk's criterion for property (T), but they meet our criterion given in corollary 3.7. Another interesting thing about the examples below is that two of them are connected to sporadic simple groups. The reader should note that connection of the second example to one of the Fischer's group isn't straightforward - the Fischer's group does not act on the GAB but is related to it . In the table below the first column indicates the Coxeter diagram (with (s, t) written above every link), the second column indicates the group associated with the GAB (if such exists), the third column indicates the universal cover and the last column indicates the reference from which the example was taken.

Table 1 - GAB's which meet out criterion

diagram	group	universal cover	reference
	Lyons	$\tilde{G}_2$	[Kan81], 4th example
	Fischer	$\tilde{C}_2$	[Kan81], 2th example
	-	$\tilde{C}_2$	[Ros86, C.6.10]

We will show that the first of those examples meets our criterion (checking the other two examples is left to the reader). In the first example, there are three types of links:

1. Bipartite graph in which $\lambda = 1$ and therefore $\overline{\lambda}_1 = \frac{1}{2}$
2. Generalized 3-gon with $s = 5$ in which $\overline{\lambda}_2 = \frac{1}{2} - \frac{\sqrt{5}}{6}$

3. Generalized 6-gon with $s = t = 5$ in which $\overline{\lambda}_3 = \frac{1}{2} - \dfrac{\sqrt{15}}{6}$

Therefore

$$\overline{\lambda}_2 + \overline{\lambda}_3 = 1 - \frac{\sqrt{5} + \sqrt{15}}{6} < 0$$

so Żuk's criterion does not hold, but

$$\overline{\lambda}_1 + \overline{\lambda}_2 + \overline{\lambda}_3 = \frac{3}{2} - \frac{\sqrt{5} + \sqrt{15}}{6} > 0$$

and

$$(\overline{\lambda}_1 + \overline{\lambda}_2)(\overline{\lambda}_1 + \overline{\lambda}_3) + (\overline{\lambda}_1 + \overline{\lambda}_2)(\overline{\lambda}_2 + \overline{\lambda}_3) + (\overline{\lambda}_1 + \overline{\lambda}_3)(\overline{\lambda}_2 + \overline{\lambda}_3) =$$

$$= 3\frac{5}{9} + \frac{15\sqrt{3}}{36} - \frac{4}{6}(\sqrt{5} + \sqrt{15}) > 0$$

therefore the criterion stated in the corollary holds.

Note that since this example was constructed using the Lyons group which act on the GAB, there is a natural extension of the Lyons group by the fundamental group of the GAB and this extension also has property (T) (either because it acts on the universal cover or becuase it is an extension of a finite group by a group with property (T)).

5.2 Hyperbolic buildings

A Dynkin diagram is of compact hyperbolic type, if it is not of finite or affine type, but every proper subdiagram is finite. For every compact hyperbolic Dynkin diagram and every finite field $\mathbb{F}_q$, Tits [Tit87] constructed a Kac-Moody group, acting cocompactly on an hyperbolic building with thickness $q + 1$. Compact hyperbolic Dynkin diagram were classified in [CCC$^+$10] but we will deal only in the 2-dimensional case, i.e. the Dynkin diagram has 3 vertices. Since the links are again generalized polygons, we can use the results of [FH64] again for the smallest positive eigenvalue of the Laplacian of each link. The table below compares the minimal q needed to assure property (T) using our Theorem 1 and [Żuk96, Theorem 1] (cases in which both theorems give the same q were omitted).

Table 2 - comparison of criteria for 2 dimensional hyperbolic buildings

diagram	minimal q by Theorem 1	minimal q by [Żuk96, Theorem 1]
	7	8
	9	11
	5	8
	7	11

References

[BdlHV08] Bachir Bekka, Pierre de la Harpe, and Alain Valette. *Kazhdan's property (T)*, volume 11 of *New Mathematical Monographs*. Cambridge University Press, Cambridge, 2008.

[BŚ97] W. Ballmann and J. Świątkowski. On L^2-cohomology and property (T) for automorphism groups of polyhedral cell complexes. *Geom. Funct. Anal.*, 7(4):615–645, 1997.

[CCC$^+$10] Lisa Carbone, Sjuvon Chung, Leigh Cobbs, Robert McRae, Debajyoti Nandi, Yusra Naqvi, and Diego Penta. Classification of hyperbolic Dynkin diagrams, root lengths and Weyl group orbits. *J. Phys. A*, 43(15):155209, 30, 2010.

[DJ00] Jan Dymara and Tadeusz Januszkiewicz. New Kazhdan groups. *Geom. Dedicata*, 80(1-3):311–317, 2000.

[FH64] Walter Feit and Graham Higman. The nonexistence of certain generalized polygons. *J. Algebra*, 1:114–131, 1964.

[Kan81] William M. Kantor. Some geometries that are almost buildings. *European J. Combin.*, 2(3):239–247, 1981.

[Kas11] Martin Kassabov. Subspace arrangements and property T. *Groups Geom. Dyn.*, 5(2):445–477, 2011.

[Ros86] L. A. Rosati, editor. *Buildings and the geometry of diagrams*, volume 1181 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1986. Lectures given at the third 1984 session of the Centro Internazionale Matematico Estivo (CIME) held in Como, August 26–September 4, 1984.

[Sha00] Yehuda Shalom. Rigidity of commensurators and irreducible lattices. *Invent. Math.*, 141(1):1–54, 2000.

[Tit81] J. Tits. A local approach to buildings. In *The geometric vein*, pages 519–547. Springer, New York, 1981.

[Tit87] Jacques Tits. Uniqueness and presentation of Kac-Moody groups over fields. *J. Algebra*, 105(2):542–573, 1987.

[Żuk96] Andrzej Żuk. La propriété (T) de Kazhdan pour les groupes agissant sur les polyèdres. *C. R. Acad. Sci. Paris Sér. I Math.*, 323(5):453–458, 1996.